

## CLIMATE


## How unusual is today's solar activity?

Arising from: S. K. Solanki, I. G. Usoskin, B. Kromer, M. Schüssler & J. Beer *Nature* 431, 1084–1087 (2004)

To put global warming into context requires knowledge about past changes in solar activity and the role of the Sun in climate change. Solanki *et al.*<sup>1</sup> propose that solar activity during recent decades was exceptionally high compared with that over the preceding 8,000 years. However, our extended analysis of the radiocarbon record reveals several periods during past centuries in which the strength of the magnetic field in the solar wind was similar to, or even higher than, that of today.

Solanki *et al.* combine radiocarbon (<sup>14</sup>C) data, visually observed sunspot numbers and models to extend the historical sunspot record over the Holocene. They exclude the most recent 100 years of the <sup>14</sup>C record, which are influenced by <sup>14</sup>C-depleted fossil-fuel emissions and atomic-bomb tests conducted since AD 1950. We extend the analysis of the radiocarbon record to AD 1950, which allows us to link the <sup>14</sup>C-based solar reconstruction to instrumental measurements of solar magnetic modulation that cover the past 68 years<sup>2,3</sup>.

The Sun influences the production rate of <sup>14</sup>C in the Earth's atmosphere by modulating the galactic cosmic-ray flux through its magnetic field. Increased magnetic field in the solar wind causes a stronger deflection of galactic cosmic rays and lower radionuclide production rates in the atmosphere and vice versa. However, the atmospheric <sup>14</sup>C concentration also depends nonlinearly on the geomagnetic field intensity and the global carbon cycle. These factors and their uncertainties need to be carefully included in the reconstruction of solar activity.



**Figure 1 | Measured atmospheric <sup>14</sup>CO<sub>2</sub>, <sup>12</sup>CO<sub>2</sub> and <sup>13</sup>CO<sub>2</sub>:<sup>12</sup>CO<sub>2</sub> ratios and model results for  $\delta^{13}\text{C}$  from AD 1500 to 1950.**  $\Delta^{14}\text{C}$  and  $\delta^{13}\text{C}$  denote the per mil deviations from a standard. **a**, Tropospheric radiocarbon from tree-ring measurements from the Northern Hemisphere<sup>4</sup> (black) and Southern Hemisphere<sup>5</sup> (brown). Error bars show one standard deviation. **b**, The Law Dome (Antarctica)  $\delta^{13}\text{C}$  record<sup>14</sup> (dots) compared with the output of the Bern model (black line). Error bars show one standard deviation.


We reconstructed <sup>14</sup>C production rates from radiocarbon records<sup>4,5</sup> (Fig. 1a) using two box-diffusion type carbon-cycle models, the box-diffusion<sup>6</sup> and the Bern model<sup>7</sup>, and a three-dimensional ocean model coupled to a four-box land biosphere. The <sup>14</sup>C production record was transformed<sup>8,9</sup> into a record of the solar modulation parameter that describes the solar influence on galactic cosmic-ray deflection by normalizing it to neutron-monitor and ionization-chamber data covering the recent decades<sup>2</sup>. Alternatively, balloon-borne estimates of galactic cosmic-ray deflection<sup>3</sup> were used instead of the ionization-chamber data. Like Solanki *et al.*<sup>1</sup>, we assumed that natural variations in the carbon cycle were small during the past millennium, which is consistent with ice-core CO<sub>2</sub> and <sup>13</sup>CO<sub>2</sub> data and models<sup>10</sup>. We considered uncertainties in <sup>14</sup>C data, fossil CO<sub>2</sub> emissions, the geomagnetic field and model parameters. Results are similar across the range of models and simulations. Calculations using the Northern<sup>4</sup> or Southern<sup>5</sup> Hemisphere <sup>14</sup>C record, or both, reveal that differences between the <sup>14</sup>C records do not affect our main conclusions.

It is standard practice to model the observed dilution of the atmospheric isotope ratios caused by the addition of isotopically depleted carbon from fossil and land-use sources. Emissions from fossil sources are prescribed, emissions from land-use sources are inferred from the atmospheric carbon budget, and the two-way exchange fluxes between reservoirs are simulated. The dilution of <sup>14</sup>C is governed by the same processes that affect <sup>13</sup>C. The good agreement between modelled <sup>13</sup>C and ice-core data (Fig. 1b) supports the reconstructed rate of <sup>14</sup>C production.

The comparison between the physical quantities, the <sup>14</sup>C-production rate and the solar-modulation parameter, and the visually based sunspot record reveals similarities and striking differences (Fig. 2). The 11-year solar cycle is distinct in all records. On the other hand, solar magnetic modulation was higher or comparable to today during the late eighteenth (and twelfth) century and around AD 1600, whereas sunspot numbers were highest over the recent decades. Sunspot numbers fell to zero during the Maunder Minimum (AD 1650–1700), whereas <sup>14</sup>C production and solar modulation continued to vary. The balloon-borne measurements<sup>3</sup> imply lower values for the solar modulation parameter than the ionization-chamber data<sup>2</sup>. Those data seem preferable because they agree better with the solar-modulation changes inferred by Solanki *et al.*<sup>1</sup> than do the balloon-borne measurements<sup>3</sup>. If

we include the Northern and Southern Hemisphere <sup>14</sup>C records (three-dimensional model) and normalize the data to the balloon-borne record, we obtain very similar levels of solar modulation (black curve in Fig. 2b).

In any case, and irrespective of the data set applied, the recent solar activity is not exceptionally high (Fig. 2). The <sup>14</sup>C results are broadly consistent with earlier reconstructions based on <sup>10</sup>Be data from the South Pole, which show that production rates around AD 1780 and in the twelfth century were comparable to those observed today<sup>11</sup>. We conclude that



**Figure 2 | Radiocarbon production rate and solar modulation parameter compared with the group sunspot number.** **a**, <sup>14</sup>C-production rates (blue) were calculated from the Northern Hemisphere <sup>14</sup>C record with two carbon-cycle models<sup>6,7</sup>; fossil-fuel emissions ( $\pm 10\%$ ) and air-sea CO<sub>2</sub> exchange rate ( $\pm 30\%$ ) were varied systematically in sensitivity calculations (various lines). The shadowed area shows the  $1\sigma$  errors resulting from 100 Monte Carlo simulations. We included Northern Hemisphere<sup>4</sup> and Southern Hemisphere<sup>5</sup> <sup>14</sup>C data in a three-dimensional ocean model calculation (green line). The black curves show the <sup>14</sup>C-production rate corrected for the geomagnetic field intensity<sup>8,9</sup>. The red curve shows the group sunspot number<sup>15</sup>. All records are normalized and show 11-yr averages. **b**, The solar modulation parameter was calculated<sup>18</sup> from the <sup>14</sup>C-production rates and from ionization-chamber and neutron-monitor data<sup>2</sup>. The grey band shows results from annual <sup>14</sup>C data, including the error range based on the Monte Carlo calculations and uncertainties in the geomagnetic data<sup>9</sup>. The black line shows 11-yr averages (best estimate). Unfiltered (blue) and 11-yr-averaged (red) sunspot numbers<sup>15</sup> are shown for comparison. The green line shows the three-dimensional model results. The purple line depicts results using an alternative neutron-monitor record<sup>3</sup> that indicates a stronger trend from 1937 to 1950 than the ionization-chamber data<sup>2</sup>.

the link between the visually based sunspot numbers and solar-modulation parameter is neither straightforward nor yet understood, and also that solar modulation must have reached or exceeded today's magnitudes three times during the past millennium.

Uncertainties in low-frequency changes increase when reconstructions are extended from the past few centuries to the past millennia. The low-frequency Holocene  $^{14}\text{C}$  variations can largely be explained by changes in the geomagnetic field as they lie within the errors of the archaeomagnetic data set used for correction<sup>12</sup>. Some palaeomagnetic records<sup>13</sup> indicate higher geomagnetic intensities around 7000 BC, which indicate that solar activity could have been lower during this period than is suggested by Solanki *et al.*<sup>1</sup>.

What do our results mean for climate change? It is speculative to translate solar magnetic modulation quantitatively into irradiance because we do not have a clear mechanistic understanding or evidence from data. Still, records of solar magnetic modulation

proxies are often used as direct indicators of solar irradiance in climate and carbon-cycle model calculations (see ref. 10, for example). The reconstruction by Solanki *et al.* implies generally less solar forcing during the past millennium than in the second part of the twentieth century, whereas our reconstruction indicates that solar activity around AD 1150 and 1600 and in the late eighteenth century was probably comparable to the recent satellite-based observations. In any case, as noted by Solanki *et al.*, solar activity reconstructions tell us that only a minor fraction of the recent global warming can be explained by the variable Sun.

**Raimund Muscheler\***, **Fortunat Joos†**,  
**Simon A. Müller‡**, **Ian Snowball‡**

\*National Center for Atmospheric Research,  
Climate and Global Dynamics Division,  
Paleoclimatology, Boulder,  
Colorado 80305-3000, USA  
e-mail: raimund@ucar.edu

‡Climate and Environmental Physics,  
Physics Institute, University of Bern,

3012 Bern, Switzerland

‡GeoBiosphere Science Centre, Quaternary  
Sciences, Lund University, 22362 Lund, Sweden

1. Solanki, S. K., Usoskin, I. G., Kromer, B., Schüssler, M. & Beer, J. *Nature* **431**, 1084–1087 (2004).
2. Beer, J. *Space Sci. Rev.* **93**, 107–119 (2000).
3. McCracken, K. G. & Heikkilä, B. in *Proc. 28th Int. Cosmic Ray Conf. 2003, Tsukuba, Japan* 4117–4120 (Univ. Acad. Press, Tokyo, 2003).
4. Stuiver, M., Reimer, P. J. & Braziunas, T. F. *Radiocarbon* **40**, 1127–1151 (1998).
5. McCormac, F. G. *et al. Radiocarbon* **44**, 641–651 (2002).
6. Siegenthaler, U. *J. Geophys. Res.* **88**, 3599–3608 (1983).
7. Joos, F. *et al. Tellus B* **48**, 397–417 (1996).
8. Masarik, J. & Beer, J. *J. Geophys. Res.* **104**, 12099–12111 (1999).
9. Yang, S., Odah, H. & Shaw, J. *Geophys. J. Int.* **140**, 158–162 (2000).
10. Gerber, S. *et al. Clim. Dynam.* **20**, 281–299 (2003).
11. Bard, E., Raisbeck, G. M., Yiou, F. & Jouzel, J. *Tellus B* **52**, 985–992 (2000).
12. Muscheler, R., Beer, J., Kubik, P. W. & Synal, H.-A. *Quat. Sci. Rev.* **24**, 1849–1860 (2005).
13. Snowball, I. & Sandgren, P. *Earth Planet. Sci. Lett.* **227**, 361–376 (2004).
14. Francey, R. J. *et al. Tellus B* **51**, 170–193 (1999).
15. Hoyt, D. V. & Schatten, K. H. *Solar Phys.* **179**, 189–219 (1998).

doi:10.1038/nature04045